

Glue drops inside hadrons

B.Z. Kopeliovich^{ab}, B. Povh^c, Ivan Schmidt^a

^aUniversidad Tecnica Federico Santa Maria, Valparaiso, Chile

^bInstitut für Theoretische Physik der Universität, Heidelberg, Germany

^cMax-Planck-Institut für Kernphysik, 69029 Heidelberg, Germany

We present experimental evidences for the existence of a semi-hard scale in light hadrons. This includes the suppression of gluon radiation that is seen in high mass hadron diffraction; the weak energy dependence of hadronic total cross sections; the small value of the Pomeron trajectory slope measured in photoproduction of J/Ψ ; the weakness of gluon shadowing in nuclei; shortage of gluons in the proton revealed by an unusual behavior of the proton structure function in the soft limit, and the enhanced intrinsic transverse momentum of quarks and gluons, which considerably exceeds the inverse hadronic size. All these observations suggest that gluons in hadrons are located within spots of a small size relative to the confinement radius.

1. Introduction

There is growing theoretical and experimental support leading towards the existence of a non-perturbative scale smaller than the usual $1/\Lambda_{QCD} \sim 1$ fm, and which is related to the gluonic degrees of freedom. First, an analysis of hadronic matrix elements of the gluonic contribution to the energy momentum tensor, using the QCD sum rules approach, gives a value of 0.3 fm for the radius of the corresponding form factor [1]. From the lattice side, numerical simulations of the gluon two point correlation function turn out a value of also 0.2 – 0.3 fm for the correlation length [2], and the energy of the QCD string appears concentrated in a tube of radius 0.3 fm in the transverse direction [3]. On the other side, it has been shown that the instanton radius peaks approximately at 1/3 fm [4]. Furthermore, high statistics data for diffractive gluon bremsstrahlung in hadronic collisions is difficult to explain unless gluons in the proton have transverse momenta as high as about 0.7 GeV [5]. This has been confirmed by studies of diffractive parton distributions, which concluded that they have a rather small transverse size [6]. What actually happens is that the smallness of the gluon clouds slows down Gribov diffusion of the gluons in transverse plane, and this results in a small slope of the Pomeron trajectory in hard reactions [7], in agreement with data. More arguments in favor of small gluonic spots coming from DIS can be found in [8]. Some of these results have been corroborated by recent studies of the spatial distribution of gluons in the transverse direction at small x [9]. Here we overview the available experimental evidences for the presence of a semihard scale in hadronic structure.

2. Why gluon radiation is suppressed

If gluons in hadrons are located within small spots of radius r_0 , they have enlarged transverse momenta $q_T \sim 1/r_0$. Such gluons cannot be resolved by soft interactions and be shaken off, which means that the bremsstrahlung cross section should be suppressed compared to perturbative estimates.

However, in the case of soft inelastic collisions followed by multiparticle production the events with or without gluon radiation look alike. In both cases the produced particles build a plateau in rapidity, and then it is difficult to find any definite signature of the radiated gluons.

Diffraction offers an exclusive possibility to identify gluon radiation. A high-energy hadron can dissociate diffractively either via excitation of the valence quark skeleton, or by radiating gluons. These two mechanisms are characterized by different dependence on the effective mass, M_X , of the excitation,

$$\frac{d\sigma(hp \rightarrow Xp)}{dM_X^2} = \begin{cases} \frac{1}{M_X^3} & \text{excitation of the quark skeleton} \\ \frac{1}{M_X^2} & \text{diffractive gluon bremsstrahlung} \end{cases} \quad (1)$$

The M_X -dependence at large M_X correlates with the spin of the slowest particle produced in the excitation. Only a vector particle, i.e. a gluon, can provide the $1/M_X^2$ dependence.

Thus, one can single out the cross section of diffractive gluon radiation from the large mass tail of the M_X -distribution. An analysis [10] of diffractive data shows that gluon radiation is amazingly weak. In order to understand that one can interpret diffraction in terms of the Pomeron-proton total cross section, as is shown in Fig. 1. If we treat the




Figure 1. The cross section of diffractive excitation of a proton expressed in terms of the total Pomeron-proton cross section.

Pomeron like a gluonic dipole, one may expect a cross section 9/4 times larger than for a $\bar{q}q$ dipole. Comparing with the pion-proton cross section, say 25 mb, one arrives at the estimate of about 50 mb. However, data suggest quite a smaller value, about 2 mb. A straightforward explanation for such a dramatic disagreement would be a much smaller size of the gluonic dipole (Pomeron) compared to the quark-antiquark dipole (pion). Thus, one concludes that gluons should be located within small spots in the proton.

Although it is not quite rigorous, one might try to estimate the diffractive radiation cross section perturbatively, and in this case the result exceeds data by more than an order

of magnitude. To reduce the cross section down to the observed value one should assume that the mean quark-gluon separation is as small as $r_0 = 0.3 \text{ fm}$ [5]. With such a modified quark-gluon light-cone distribution function, the effective triple-Pomeron coupling has the form [5],

$$G_{3\mathbf{P}}(0) \equiv (1 - x_F)^{\alpha_{\mathbf{P}}(0)} \left. \frac{d\sigma_{sd}(pp \rightarrow pX)}{dx_F dp_T^2} \right|_{p_T=0} = \frac{81\alpha_s\sigma_0}{(16\pi)^2} \ln \left[\frac{2(r_0^2 + R_0^2)^2}{R_0^2(2r_0^2 + R_0^2)} \right]. \quad (2)$$

Here we assume that $1 \gg 1 - x_F \gg s_0/s$, where $s_0 \sim 1 \text{ GeV}^2$. The energy dependent parameters $\sigma_0(s)$ and $R_0(s)$ controlling the shape of the universal dipole cross section [11] are defined in [5]. With $r_0 = 0.3 \text{ fm}$, the triple-Pomeron coupling eq. (2) agrees with the result of the triple-Regge analysis [10] of single diffractive data.

3. Why energy dependence of hadronic cross sections is so weak

It is well known that hadronic cross sections rise with energy approximately as s^ϵ , where the exponent is quite small, $\epsilon \approx 0.1$. What is the origin of this small number? We do not expect any small parameters in the soft regime of strong interactions.

This problem is closely related to the topic of the previous section. In fact, the energy dependence is driven by gluon radiation which turns out to be suppressed. Before we saw a manifestation of this effect in diffraction, now in the total inelastic cross section.

Without gluon radiation the geometric cross section of two hadrons would be constant, since their transverse size is Lorentz invariant, i.e. is energy independent. The phase space for one gluon radiation is proportional to $\ln(s)$, so multigluon radiation leads to powers of $\ln(s)$ in the cross section. The calculations performed in [12] confirm this. The hadronic cross section was found to have the following structure,

$$\sigma_{tot} = \sigma_0 + \sigma_1 \left(\frac{s}{s_0} \right)^\Delta, \quad (3)$$

where σ_0 is the energy independent term related to hadronic collisions without gluon radiation. The second term in (3) is the contribution of gluon bremsstrahlung to the total cross section. Here the parameter σ_1 is expected to be small due to the smallness of the gluonic spots. Indeed, it was found in [12] that $\sigma_1 = 27 C r_0^2/4$, where factor $C \approx 2.4$ is related to the behavior of the dipole-proton cross section, calculated in Born approximation at small separations, $\sigma(r_T) = C r_T^2$ at $r_T \rightarrow 0$.

The energy dependence of the second term in (3) was found to be rather steep, $\Delta = 4\alpha_s/3\pi = 0.17$. This exponent seems to be too large compared to the experimentally measured $\epsilon \approx 0.1$. There is, however, no contradiction due to the presence of the large energy independent term in (3). Approximating the cross section (3) by a simple power dependence on energy, the effective exponent reads,

$$\epsilon = \frac{\Delta}{1 + \sigma_0/\sigma_1 (s/s_0)^{-\Delta}} \quad (4)$$

So, one should expect a growing steepness of the energy dependence for the total cross section. One can estimate the value of r_0 demanding the effective exponent to be $\epsilon \approx 0.1$

in the energy range of fixed target experiments, say at $s \sim 1000 \text{ GeV}^2$. With $\sigma_0 = 40 \text{ mb}$ found in [12] one gets $r_0 = 0.3 \text{ fm}$.

Thus, the observed slow rise of the total hadronic cross sections provides another evidence for the existence of small gluonic spots with transverse size $r_0 \sim 0.3 \text{ fm}$.

One may expect a steeper energy dependence for heavier flavors. Indeed, for J/Ψ -proton scattering σ_0 is so small, that $\epsilon \approx \Delta$. Indeed, data for J/Ψ photoproduction from HERA [13] show that $\epsilon \approx 0.2$. One should be careful, however, interpreting the data within the vector dominance model [14], and remember that Eq. (3) was derived assuming that r_0 is much smaller than the hadronic size, otherwise interferences should be included.

4. Why diffractive cone shrinks with energy slowly

The prediction of a shrinkage of the diffraction cone has been one of the first achievements of the Regge theory. Indeed, data show that the elastic slope in hadronic collisions rises with energy as, $B_{el}(s) = B_0 + 2\alpha'_{\mathbf{P}} \ln(s/s_0)$, where $\alpha'_{\mathbf{P}} \approx 0.25 \text{ GeV}^{-2}$. This is about four times smaller than in binary processes mediated by other Reggeons, $\alpha'_{\mathbf{R}} \approx 1 \text{ GeV}^{-2}$. Why?

The diffractive cone shrinkage is usually related to Gribov diffusion of gluons in the transverse plane. If each "step" in impact parameters, $\Delta b^2 = r_0^2$, is small, the diffusion should proceed slowly. Indeed, a rather small value of the Pomeron trajectory slope was predicted in [12],

$$\alpha'_{\mathbf{P}} = \frac{1}{2} \frac{dB_{el}}{d \ln(s/s_0)} = \frac{\alpha_s}{3\pi} r_0^2 = 0.1 \text{ GeV}^{-2}. \quad (5)$$

This seems to be too small, in strong contradiction with value 0.25 GeV^{-2} known from data for the elastic slope. One may wonder, why the same model [12] which predicts (5) describes well data for elastic slope, as is demonstrated in Fig. 2? The relatively large value of $\alpha'_{\mathbf{P}}$, turns out to result from unitarity saturation. Indeed, the elastic differential cross section (actually the amplitude) Fourier transformed to impact parameter representation (see details in [12]) demonstrate unitarity saturation at small impact parameters. In spite of the observed rise of the total cross section with energy, there is no room for further growth at small impact parameters, only the amplitudes of peripheral collisions rise with energy. This leads to a rising with energy radius of interaction directly related to the elastic slope. Thus, a substantial part of the observed energy dependence of the elastic slope and of the effective $\alpha'_{eff} = 0.25 \text{ GeV}^{-2}$ is related to saturation of the unitarity bound. How to disentangle the two effects?

To get rid of unitarity corrections one can consider the interaction of a small dipole with a proton. For example photoproduction of a heavy quarkonium [7], or high Q^2 electroproduction of ρ . Then, the elastic amplitude is too small to be affected by unitarity (absorptive) corrections, and the energy dependence of the slope must be solely due to the rise of the gluon clouds, i.e. Gribov diffusion. This expectation of [12] was nicely confirmed in elastic photoproduction $\gamma + p \rightarrow J/\Psi + p$ measured by the ZEUS experiment [13] which found $\alpha'_{\mathbf{P}} = 0.115 \pm 0.018 \text{ GeV}^{-2}$. The data and fit are depicted in Fig. 3




Figure 2. Elastic slope for $pp/\bar{p}p$ (bottom/upper curves) collisions, calculated in [12] with the small Pomeron slope according to (5). The references to the data can be found in [12].




Figure 3. Elastic slope in exclusive photoproduction $\gamma + p \rightarrow J/\Psi + p$. Data are from the ZEUS experiment [13]. The solid curve is a fit with $\alpha'_{\mathbf{P}} = 0.115 \pm 0.018 \text{ GeV}^{-2}$, in good agreement with (5).

5. Why gluon shadowing is so weak

Although bound nucleons in a nucleus are rather well separated, Lorentz-boosted into the infinite momentum frame they start overlapping at small Bjorken x . This happens since Lorentz contraction of the gluonic clouds at small x is much weaker than at large x . If the clouds originated from different nucleons overlap, they may fuse reducing the gluon density at small x . This phenomenon is called gluon shadowing. At first glance this might be a considerable effect, since gluons interact stronger than quarks. However, one should be cautious, since a similar naive expectation of a large cross section for gluonic dipoles failed when was confronted with data (see Sect. 2), and both phenomena have common roots.

Even if gluonic clouds overlap in the longitudinal direction, one should make sure that it also happens in the transverse plane. This is not obvious if the clouds are small. In a nucleus the mean number of gluonic spots overlapping at a given impact parameter is,

$$\langle n \rangle = \frac{3\pi}{4} r_0^2 \langle T_A \rangle, \quad (6)$$

where $\langle T_A \rangle \approx \rho_A \frac{4}{3} R_A$ is the mean nuclear thickness; $\rho_A \approx 0.16 \text{ fm}^{-3}$ is the nuclear density; and $R_A \approx 1.14 \text{ fm} \times A^{1/3}$ is the nuclear radius. Notice that r_0 is the mean diameter of a dipole, rather than radius, which explains the factor $1/4$ in (6).

According to (6) even for the heaviest nuclei only very few gluonic clouds have a chance to overlap at a given impact parameter, $\langle n \rangle \approx 0.3$. Therefore, the longitudinal overlap of gluons originating from different nucleons does not lead to gluon interaction if the transverse overlap is so small. This may substantially weaken all collective phenomena at small x including color glass condensate [15].

Thus, another consequence of smallness of the gluonic spots is a very weak gluon shadowing, predicted in [5]. Fig. 4 demonstrates results of calculations performed in [5] by different methods and at different scales $Q^2 \approx 1/r_0^2$ and 4 GeV^2 . Even at x as low as 10^{-4} shadowing corrections for medium heavy nuclei are only 10%.




Figure 4. Predictions for gluon shadowing based on the small gluonic spot structure [5]. The two sets of curves are calculated by different methods and at different scales, at $Q^2 \approx 1/r_0^2$ (thin curves) and at $Q^2 = 4 \text{ GeV}^2$ (thick curves). Each set is calculated for carbon, iron and lead (from top to bottom).




Figure 5. Gluon shadowing in calcium suggested by DGLAP analysis of DIS data on nuclei. The solid curve is from an analysis [18] done in the next-to-leading order approximation. The dashed curve is the result of a leading order analysis [19].

Unfortunately, the gluon distribution function is difficult to measure, since the main partonometer, deep-inelastic scattering (DIS), probes directly only quarks and antiquarks. The gluon distribution can be accessed only via Q^2 evolution. Although previous attempts to single out the gluon distribution in nuclei from DIS resulted in big uncertainties, a recent next-to-leading order analysis [18] turned out to be quite sensitive to gluons. Gluon shadowing extracted from data is depicted in Fig. 5 for calcium [18]. Dashed curve shows the results of a LO analysis [19]. We do not compare with the gluon shadowing suggested in [20], since it was ad hoc. The results of the DGLAP analyses agree quite well with predictions [5] within the theoretical and fitting uncertainties. Other gluon shadowing calculations missing the effect of small gluonic spots dramatically overestimate the magnitude of gluon shadowing.

Suppression of gluon radiation due to smallness of gluonic spots also leads to a substantial reduction of the effect of color glass condensate. This is demonstrated in Fig. 6

where p_T -dependent nuclear ratios are calculated for long-range ($r_0 \sim 1/\Lambda_{QCD}$) and short range ($r_0 \sim 0.3$ fm) gluons [16]. Gluonic spot structure of the proton also leads to a very weak Cronin enhancement predicted for RHIC (confirmed) and LHC [17].





Figure 6. Normalized ratios of gluon radiation cross sections in quark-nucleus over quark-nucleon collisions. The ratios are plotted as function of gluon transverse momentum. Calculations [16] are performed for lead in two cases of long-range and short range ($r_0 = 0.3$ fm) gluons. In the latter case the magnitude of the color glass condensate effect is much smaller.



resolution one can see gluons whose Fermi motion is much more intensive, since they are confined within small spots, $\langle k_T \rangle \sim 1/r_0 \approx 700$ MeV. There should be manifestations of such an enhanced Fermi motion in reactions and observables sensitive to the primordial parton momentum.

7.1. Sea-Gull effect

The projectile quark and gluons lose (at least partially) coherence and emerge from the interaction area as forward jets. These jets having rather small transverse momenta hardly can be reconstructed. However, inclusively detected hadrons carrying fraction z of the initial quark/gluon correspondingly have same fraction of the jet transverse momentum. Therefore the mean transverse momentum of inclusive hadrons should rise as function of Feynman x_F . This effect is indeed observed in data [22] and it allows to measure the jet transverse momentum which results from the momentum transfer gained in the interaction and from the primordial intrinsic momentum of the parton. The latter can be extracted from data and it turns out to be rather high, $0.5 - 1$ GeV, the value anticipated in the model of gluonic spots.

7.2. Unintegrated gluon distribution

Direct information about transverse momenta of gluons in a hadron comes from the unintegrated distribution function. There are few phenomenological distributions in the literature. A quite popular one [23] employs the saturated shape of the dipole cross section fitted to HERA data for the proton structure function at small Bjorken x ,

$$\mathcal{F}(x, k_T^2) = \frac{3 \sigma_0 R_0^2(x) k_T^4}{16 \pi^2 \alpha_s(k_T^2)} \exp\left[-\frac{1}{4} R_0^2(x) k_T^2\right], \quad (7)$$

where $R_0(x) = 0.4 \text{ fm} \times (x/x_0)^{0.144}$; $x_0 = 0.0003$. From this distribution the mean transverse momentum squared is $\langle k_T^2 \rangle = 12/R_0^2(x)$. For instance, at $x = 0.01$ this is $\sqrt{\langle k_T^2 \rangle} \approx 1$ GeV. Although the distribution Eq. (7) falls off steeply at large k_T , this does not have a big effect on its mean value. Notice that for the small k_T region an energy dependent saturated shape of the dipole cross section is more appropriate [5], which results in a smaller value for the mean transverse momentum of gluons, $\sqrt{\langle k_T^2 \rangle} \approx 0.6$ GeV [24].

Both estimates of the mean transverse momentum are quite large indicating at localization of gluons within small spots.

7.3. σ_L/σ_T in DIS

Of course, not only gluons get increased transverse momenta, but also valence quarks if they are probed with sufficient resolution, $Q^2 \gg 1/r_0^2$. The ratio of the longitudinal-to-transverse DIS cross sections, $R = \sigma_L/\sigma_T$, is sensitive to the transverse motion of valence quarks (at large x), since otherwise it vanishes in LO at high energy ν (or high Q^2 and fixed x) as $R = g(x)Q^2/\nu^2$ according to the Callan-Gross relation [25,26].

In order to incorporate the effect of transverse motion of the quarks, a parametrization $R = cQ^2/(Q^2 + d^2)^2$ was used in [27] to fit data, with $d^2 = 0.99 \pm 0.39 \text{ GeV}^2$. This result agrees with the expected $d \sim 1/r_0$.

Notice that data for R_x, Q^2 is well explained in the dipole approach [28] with a saturated cross section. This is not a surprise, since according to our discussion in the previous section the dipole cross section bears information about enhanced gluon momenta.

Acknowledgments: We are thankful to Hans-Jürgen Pirner for helpful discussions. This work was supported in part by Fondecyt (Chile) grants numbers 1030355 and 1050519, and by DFG (Germany) grant PI182/3-1.

REFERENCES

1. V.M. Braun, P. Górnicki, L. Mankiewicz, Phys. Lett. B302 (1993) 291.
2. M. D’Elia, A. Di Giacomo, Phys. Lett. B408 (1997) 315.
3. G.S. Bali, Int. Conference in Quark Confinement and Hadron Spectrum, Newport News (Virginia, USA), June 1998.
4. E.V. Shuryak, Phys. Lett. B486 (2000) 378.
5. B.Z. Kopeliovich, A. Schäfer, and A.V. Tarasov Phys. Rev. D62 (2000) 054022.
6. F. Hautmann, D.E. Soper, Phys. Rev. D63 (2000) 011501.
7. B.Z. Kopeliovich and B. Povh, J. Phys. G30 (2004) S999.
8. E. Shuryak and I. Zahed, Phys. Rev. D69 (2004) 014011.
9. M.I. Strikman and C. Weiss, Phys. Rev. D69 (2004) 054012.
10. Yu.M. Kazarinov, B.Z. Kopeliovich, L.I. Lapidus, and I.K. Potashnikova, Sov. Phys. JETP **70** (1976) 1152.
11. B.Z. Kopeliovich, L.I. Lapidus, and A.B. Zamolodchikov, Sov. Phys. JETP Lett. **33**, 595 (1981); Pisma v Zh. Exper. Teor. Fiz. **33**, 612 (1981).
12. B.Z. Kopeliovich, I.K. Potashnikova, B. Povh, and E. Predazzi, Phys. Rev. Lett. 85 (2000) 507; Phys. Rev. D63 (2001) 054001.
13. ZEUS Collaboration, S. Chekanov et al., Eur. Phys. J. C24 (2002) 345.
14. J. Hüfner and B.Z. Kopeliovich, Phys. Lett. B426 (1998) 154.
15. L. McLerran and R. Venugopalan, Phys. Rev. D49 (1994) 2233.
16. B.Z. Kopeliovich, A. Schäfer, and A.V. Tarasov, Phys. Rev. C59 (1999) 1609.
17. B.Z. Kopeliovich, J. Nemchik, A. Schäfer, and A.V. Tarasov, Phys. Rev. Lett. 88 (2002) 232303.
18. D. de Florian and R. Sassot, Phys. Rev. D69 (2004) 074028.
19. M. Hirai, S. Kumano and M. Miyama, Phys. Rev. D64 (2001) 034003.
20. K. J. Eskola, V. J. Kolhinen, and P. V. Ruuskanen, Nucl. Phys. **B535** (1998) 351.
21. ZEUS Collaboration, J. Breitweg et al., Eur. Phys. J. C7 (1999) 609.
22. NA22 Collaboration, N.M. Agababyan et al., Phys. Lett. B320 (1994) 411.
23. K. Golec-Biernat and M. Wüsthoff, Phys. Rev. D59 (1999) 014017.
24. M.B. Johnson, B.Z. Kopeliovich and A.V. Tarasov, Phys. Rev. C **63**, 035203 (2001).
25. C.G. Callan and D.J. Gross, Phys. Rev. Lett. 22 (1969) 126.
26. J.F. Gunion and R.L. Jaffe, Phys. Rev. D8 (1973) 3215.
27. A. Bodek et al., Phys. Rev. D20 (1979) 471;
28. B.Z. Kopeliovich, J. Raufeisen and A.V. Tarasov, Phys. Rev. C62 (2000) 035204.